

Effects of Pre-Drying, Delayed Cooling, and Carbon Dioxide on Skin Blackening Disorder in Asian pear (*Pyrus pyrifolia* Nakai) ‘Chuhwangbae’

Ug-Yong Lee¹, Mao-Hua Wang², Tae-Min Bae², Sung-Joo Kim², Jin-Ho Choi¹,
Young-Jik Ahn³, and Jong-Pil Chun^{2*}

¹Pear Research Institute, National Institute of Horticultural and Herbal Science, Naju 58216, Korea

²Department of Horticultural Science, Chungnam National University, Daehak-ro 99, Daejeon 34134, Korea

³Department of Horticulture & Forestry, PaiChai University, Daejeon 35345, Korea

*Corresponding author: jpchun@cnu.ac.kr

원예과학기술지 36권 3호에 게재된 논문의 사사가 누락되어 바로잡습니다.

This work was carried out with the support of “Cooperative Research Program for Agriculture Science & Technology Development (PJ01190803)” Rural Development Administration, Republic of Korea.

Received: October 31, 2017

Revised: December 29, 2017

Accepted: December 31, 2017

 OPEN ACCESS



HORTICULTURAL SCIENCE and TECHNOLOGY

36(6):900, 2018

URL: <http://www.kjhst.org>

pISSN : 1226-8763

eISSN : 2465-8588

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright©2018 Korean Society for Horticultural Science.